

Climate commitment in an uncertain world

K. C. Armour¹ and G. H. Roe²

Received 18 October 2010; revised 28 November 2010; accepted 2 December 2010; published 15 January 2011.

[1] Climate commitment—the warming that would still occur given no further human influence—is a fundamental metric for both science and policy. It informs us of the minimum climate change we face and, moreover, depends only on our knowledge of the natural climate system. Studies of the climate commitment due to CO₂ find that global temperature would remain near current levels, or even decrease slightly, in the millennium following the cessation of emissions. However, this result overlooks the important role of the non-CO₂ greenhouse gases and aerosols. This paper shows that global energetics require an immediate and significant warming following the cessation of emissions as aerosols are quickly washed from the atmosphere, and the large uncertainty in current aerosol radiative forcing implies a large uncertainty in the climate commitment. Fundamental constraints preclude Earth returning to pre-industrial temperatures for the indefinite future. These same constraints mean that observations are currently unable to eliminate the possibility that we are already beyond the point where the ultimate warming will exceed dangerous levels. Models produce a narrower range of climate commitment, but under-sample observed forcing constraints. **Citation:** Armour, K. C., and G. H. Roe (2011), Climate commitment in an uncertain world, *Geophys. Res. Lett.*, 38, L01707, doi:10.1029/2010GL045850.

1. Introduction

[2] Our ability to predict future climate changes rests fundamentally on two factors: firstly, how our future human activities will influence climate forcing and secondly, how our models of the climate system translate that forcing into climate change. The first factor depends on societal choices beyond the scope of science. The second factor depends on our confidence in the climate models. In turn, this confidence is predicated on the ability of the models to reproduce past climate changes, given our knowledge of previous human (and other) influences.

[3] The concept of a ‘climate commitment’—the climate change that would still occur given no further human influence—has proven useful in distinguishing between these two factors of climate prediction. It allows for a clear separation between the uncertainties in our physical climate models, which we wish to study, and the highly-uncertain future human influence on climate. The climate commitment can also be regarded as the minimum climate change we are consigned to because of human activities already undertaken.

¹Department of Physics, University of Washington, Seattle, Washington, USA.

²Department of Earth and Space Sciences, University of Washington, Seattle, Washington, USA.

[4] Early efforts to estimate climate commitment considered the additional warming that occurs as the climate system comes into equilibrium with the present atmospheric composition and radiative forcing. Under this assumption, an additional warming of about 0.6°C is ‘in the pipeline’ due to the thermal inertia of the world oceans [Wigley, 2005; Meehl *et al.*, 2005; Hansen *et al.*, 2005], committing us to future climate change that approaches ‘dangerous’ levels [Ramanathan and Feng, 2008].

[5] There has recently been a resurgence of interest in the climate commitment [Ramanathan and Feng, 2008; Hare and Meinshausen, 2006; Plattner *et al.*, 2008; Solomon *et al.*, 2009; Matthews and Weaver, 2010] in which an alternative, ‘zero emissions’, definition has been proposed. Under zero emissions, the atmospheric composition changes according to natural processes, and future warming is determined by only the physical inertia of the climate system and the residual greenhouse gas climate forcing. Matthews and Weaver [2010] argue that this definition is the correct measure of the present climate commitment. They make the worthwhile and important point that the previous measure—constant climate forcing—conflates the physical response of the climate system to past emissions with the response to the future emissions that are necessary to maintain a constant atmospheric composition.

[6] Several studies [Plattner *et al.*, 2008; Solomon *et al.*, 2009; Matthews and Weaver, 2010] consider the zero emissions commitment with respect to CO₂. Carbon dioxide is naturally removed from the atmosphere on multiple time scales. Under zero emissions, CO₂ would fall off to about 40% of its peak enhancement above pre-industrial levels within a few centuries [Solomon *et al.*, 2009], while full recovery would occur over hundreds of thousands of years [Archer, 2005]. Effectively then, this residual 40% defines the ultimate radiative forcing ($\equiv R_{\infty}$) with which the climate must come into equilibrium. In such a zero emissions scenario, global average surface temperature is projected to remain near current levels, or even decrease slightly, in the millennium following the cessation of emissions [Plattner *et al.*, 2008; Solomon *et al.*, 2009; Matthews and Weaver, 2010]. However, these studies have overlooked the important role of the non-CO₂ greenhouse gases (such as methane and nitrous oxide) and aerosols. Aerosols are widely known to be one of the chief uncertainties in the modern climate, and make a considerable difference to the answer. Ramanathan and Feng [2008] do consider the effect of removing anthropogenic aerosols, however they fix CO₂ at modern levels. The full consequences of the cessation of human activities must include both influences.

2. Transient and Ultimate Climate Commitment

[7] Following the elimination of emissions, aerosols would fall to their pre-industrial levels on time scales of days to

weeks [Forster *et al.*, 2007], while the non-CO₂ greenhouse gases would persist for decades to centuries [Forster *et al.*, 2007; Solomon *et al.*, 2009]. The sudden loss of the cooling effect of aerosols would result in a rapid transient warming as the surface temperature adjusts to the full greenhouse gas radiative forcing. Due to this significant transient warming, we propose two separate measures of climate commitment: a ‘transient commitment’, defined by the peak temperature following the cessation of emissions; and an ‘ultimate commitment’, defined by the temperature once the climate system has fully equilibrated with the persistent fraction of the CO₂ radiative forcing.

[8] How well constrained is the climate commitment? Conservation of energy must obviously apply to the global energy budget, a linearization of which is

$$H = R - \lambda^{-1}T, \quad (1)$$

where λ is the climate sensitivity parameter, T is the global average surface temperature (above pre-industrial), R is the radiative forcing, and H is the ocean heat uptake.

[9] For a permanent forcing R_∞ , H must ultimately go to zero giving an ultimate commitment of

$$T_\infty = \lambda R_\infty. \quad (2)$$

Eliminating λ gives

$$T_\infty = \left(\frac{R_\infty}{R - H} T \right). \quad (3)$$

Thus, T_∞ depends only on observed constraints (T , H , and R) and the ultimate forcing (R_∞). For the current climate, T is $0.76 \pm 0.11^\circ\text{C}$ (1σ) [Trenberth *et al.*, 2007] and H is $0.74 \pm 0.08 \text{ W m}^{-2}$ (1σ) [Lyman *et al.*, 2010; Purkey and Johnson, 2010].

[10] The Intergovernmental Panel on Climate Change Fourth Assessment Report (IPCC AR4) outlines constraints on R . Total anthropogenic radiative forcing is approximately 1.6 W m^{-2} , with a 90% confidence range of 0.6 W m^{-2} to 2.4 W m^{-2} [Forster *et al.*, 2007]. Therefore, T and H are well constrained, R less so. Uncertainties in aerosols (-0.5 W m^{-2} to -2.2 W m^{-2}) [Forster *et al.*, 2007] dominate the uncertainty in R , and thus dominate the uncertainty in T_∞ .

[11] The time evolution of climate requires a representation of the ocean, for which we use a simple upwelling-diffusion model. The model is the same as that of Baker and Roe [2009], which is similar in form to those used in previous studies [e.g., Hoffert *et al.*, 1980; Raper *et al.*, 2001]. All parameters are as described by Baker and Roe [2009], except R and λ , which we vary as described below. Such models are robust, and successfully reproduce observations of ocean heat uptake at the global scale [Raper *et al.*, 2001].

3. Results

[12] The weak bounds on aerosols means a broad envelope of uncertainty in total forcing over the industrial era. This is illustrated in Figure 1a, where an idealized representation of forcing trends has been employed. Forcing reaches its modern value in year 200, and from then on a climate commitment scenario is assumed. Once emissions

are terminated, R is governed by the respective decay time scales of the various atmospheric constituents (see Figure 1).

[13] From equation (1) the relatively strong constraints on T and H mean that R and λ can be thought of as pairs wherein strong (weak) aerosol forcing is balanced by high (low) climate sensitivity. This compensation occurs within AR4 and older models [Schwartz *et al.*, 2007; Kiehl, 2007; Knutti, 2008]. Figure 1b shows temperature trajectories for pairs of R and λ , whereby past temperature trends are approximately reproduced. It is a graphical representation of the inherent trade-off between uncertainties in climate forcing and uncertainties in global temperature following the cessation of emissions: even though past temperature changes are well constrained and future forcing (under zero emissions) well understood, uncertainty in past forcing implies uncertainty in future temperatures.

[14] It is important to emphasize that R and λ are not independent. In other words, a high climate sensitivity and a low aerosol forcing are inconsistent with the observed constraints on surface temperature and ocean heat uptake. Two recent studies that consider the effects of the loss of aerosols [Hare and Meinshausen, 2006; Ramanathan and Feng, 2008] treat R and λ as independent, and also fail to span the full range of either R or λ . This has the effect of producing a narrower range of climate commitment than allowed by propagating the observed constraints through equation (3).

[15] We next reproduce and explain the results of previous studies [Plattner *et al.*, 2008; Solomon *et al.*, 2009; Matthews and Weaver, 2010] that considered climate commitment with respect to only CO₂ emissions (non-CO₂ greenhouse gases and aerosols remain at their modern concentrations). For modal estimates of modern radiative forcing, this gives $R_\infty = 0.8 \text{ W m}^{-2}$. The dashed black lines in Figure 1 show this forcing and the response—a gentle decline in temperature following the cessation of CO₂ emissions. The result follows directly from surface energetics (equation (3)): R_∞ is very near the modal value of the current surface forcing ($R - H \approx 1 \text{ W m}^{-2}$) so the ratio of forcings (i.e., $R_\infty/(R - H)$) and therefore the ratio of the responses (i.e., T_∞/T) is near, but slightly less than, one.

[16] Turning now to the case in which all anthropogenic emissions cease, there is an immediate unmasking of greenhouse gas forcing as aerosols are quickly washed from the atmosphere. The effect is an abrupt rise in climate forcing (Figure 1a) to a peak value of around 2.7 W m^{-2} , which is relatively well constrained as it depends only on greenhouse gases. The response is a rapid warming (Figure 1b), with a transient commitment of up to 0.9°C above the modern temperature. Thereafter, forcing declines over the next few centuries as greenhouse gases are partially, but not completely, removed from the atmosphere. At the low end of the climate response, temperature falls to less than half of its peak value. At the high end, temperature continues to increase because the system has not yet attained equilibrium due to the long adjustment time scales of high sensitivity systems [Baker and Roe, 2009].

[17] We note that while simple upwelling-diffusive climate models, such as the one used here, are able to reproduce observed climate trends, they do not accurately capture the complexities of ocean heat uptake at the regional scale [Gregory, 2000] and likely underestimate the long-term

temperature response to forcing [Winton *et al.*, 2009]. While the details of any particular temperature trajectory are model dependent, the overall form of the temperature response is a fundamental consequence of three basic and robust climate properties: the unmasking of climate forcing by the loss of anthropogenic aerosols, the long lifetime of greenhouse gases in the atmosphere, and the thermal inertia of the ocean.

[18] The long-term temperature response depends only on modern surface energetics and R_∞ . Figure 1 accounts only for uncertainties in aerosols. This gives an ultimate commitment (above pre-industrial) of $T_\infty = 0.6^\circ\text{C}$ with a 90% confidence range of 0.3°C to 7.2°C , which follows directly from equation (3) or by integrating the climate model to equilibrium.

[19] The lower bound on climate commitment is robust due to the form of equation (3). On the other hand, the upper bound is very sensitive to uncertainties in observed global energetics and R_∞ . We do not account here for uncertainties in the biogeochemical cycle (e.g., uncertainty in the lifetimes of greenhouse gases or the residual atmospheric CO_2 concentration). Moreover, following the IPCC, we have taken the 90% confidence interval on aerosol climate forcing: if one were to factor in other sources of uncertainty, in either ocean heat uptake or greenhouse gas forcing, or use more conventional statistical bounds (i.e., a 95% range), one could not rule out the disconcerting possibility that the observed 20th century warming has transpired with little to no effective surface forcing (i.e., $R \approx H$).

4. Discussion

[20] The above analysis showed that current observational constraints allow the possibility of a very large climate commitment. Do narrower bounds exist? The ultimate commitment can alternatively be expressed as a function of λ (equation (2)), reasonable bounds on which can be inferred from IPCC AR4 in terms of a ‘likely’ (>66% probability) and ‘very likely’ (>90% probability) range for climate sensitivity [Hegerl *et al.*, 2007]. Exploiting the fundamental relationship between R and λ , and reversing the above arguments, these IPCC constraints on λ provide constraints on R (Figure 2a). Any value of λ within the IPCC range still implies a significant transient warming (Figure 2b), and there remains a substantial uncertainty in the ultimate commitment (though the range is smaller than that based on observational constraints).

[21] The ability of the IPCC AR4 fully coupled climate models (hereafter AR4 models) to reproduce 20th century surface temperature [Knutti, 2008] and ocean heat uptake [Plattner *et al.*, 2008], under substantial aerosol uncertainty, has been suggested to give a false sense of the accuracy with which future climate can be predicted [Schwartz *et al.*, 2007]. However, AR4 models have achieved consistency with the observational record, in part, through compensation between R and λ [Knutti, 2008]. As argued by Knutti [2008], such model tuning—whether explicit or implicit—is not problematic provided that we interpret models as conditional on observations. In other words, models satisfy equation (1) subject to relatively tight constraints on T and H . Accurate simulation of 20th century climate may then be viewed as a necessary, but not sufficient, condition for the ability to simulate future climate, and does not alone create overcon-




Figure 1. Observational constraints on climate forcing and temperature response. (a) Idealized representation of forcing trends. Forcing is ramped linearly to its modern value in year 200, and a zero emissions scenario is assumed thereafter. Upon zero emissions, aerosols and tropospheric ozone are specified to fall to pre-industrial levels immediately. Long-lived greenhouse gases decline at their respective (e -folding) time scales [Forster *et al.*, 2007]: 12 years for methane; 114 years for nitrous oxide; 75 years (a representative lifetime) for halocarbons. Carbon dioxide falls to 40% of its peak value (above pre-industrial) with a decay time scale of 170 years [Forster *et al.*, 2007]. Radiative forcing is calculated using the simplified expressions of Myhre *et al.* [1998]. The light blue shading is the 90% confidence interval on trajectories of R as allowed by observations, where only uncertainty in aerosols is considered. The solid black line shows the modal value of R . The dashed black line shows a scenario in which aerosols and non- CO_2 greenhouse gases are held fixed at their modern concentrations upon the elimination of CO_2 emissions. (b) As for Figure 1a, but modeled temperature response. Values of λ have been paired with values of R so that individual temperature trajectories are tightly constrained, analogous to the situation for modern observations.

fidence in model skill. Indeed, the light blue trajectories in Figure 2b clearly demonstrate the ability, with a model, to reproduce the 20th century temperature record yet still span the full range of uncertainty in climate commitment as allowed by observations.

[22] The difference between the AR4 model range of climate commitment and the range allowed by observations can instead be attributed to an inconsistency between R in models and R in observations—the range of forcing among the different AR4 models [Knutti, 2008] spans only the ‘likely’ range of forcing in Figure 2a. How can models and




Figure 2. Illustration of what IPCC bounds on climate sensitivity imply for constraints on past climate forcing and future temperature response. (a) Radiative forcing and (b) temperature response, as in Figure 1. The dark blue shading shows the IPCC AR4 ‘likely’ range of climate sensitivity (2°C to 4.5°C). The medium blue shading shows the IPCC AR4 ‘very likely’ range of climate sensitivity (1.5°C to 10°C —IPCC AR4 [Hegerl et al., 2007] truncates the probability distributions of climate sensitivity at 10°C so we take this value as representative of the upper bound on the ‘very likely’ range). For comparison, the light blue shading shows the 90% confidence interval as allowed by observations, as in Figure 1. A wedge in the lower range of possible forcing translates to a wedge in the higher range of possible temperature response.

observations be reconciled? One way would be to achieve substantially more accurate observations of the Earth’s radiative budget. In particular, emphasis should be placed on ruling out the very low values of R that correspond to very high values of committed warming.

[23] The alternative approach is to create populations of climate models that deliberately exploit tuning to fully span the uncertainty in climate forcing (and the implied range of climate sensitivity necessary to reproduce the observed temperature record), and then to demonstrate that some pairs of R and λ are inconsistent with some aspect of either the instrumental record (e.g., interannual variability, seasonal variability, spatial patterns of warming, or volcanic eruptions), or reconstructions of past climates [see Hegerl et al., 2007; Knutti and Hegerl, 2008; Edwards et al., 2007, and references therein]. Studies that pursue this approach produce a variety of distributions for climate sensitivity, many narrower than that inferred from observational constraints, some narrower than even the IPCC ‘likely’ range [Allen et al.,

2007; Hegerl et al., 2007; Edwards et al., 2007; Knutti and Hegerl, 2008]. An implication then, would be that the range of uncertainty in the climate commitment could be narrowed as well. Achieving convergence among these different distributions depends on understanding the differing assumptions and structural uncertainties in, and the interdependence of, the respective frameworks [Frame et al., 2005; Allen et al., 2007; Knutti and Hegerl, 2008; Knutti, 2010]. Arguably, an important measure of the value added by models will be when the consensus is reached that such studies provide narrower constraints on the modern climate forcing than that currently provided by direct observations. The discrepancy between the reported ranges of uncertainty in climate sensitivity and observations of aerosol forcing is an important one for future rounds of the IPCC reports to resolve. Until then, model-based estimates should be treated carefully, and probably represent an undersampling of the possible climate commitment.

5. Conclusions

[24] The results presented here depend only on three straightforward and well-understood aspects of climate: the net cooling effect of aerosols, the large spread of uncertainty in aerosol forcing (or, equivalently, climate sensitivity), and the long atmospheric lifetimes CO_2 and other greenhouse gases. In combination they lead to considerable uncertainty in the transient and ultimate climate commitments.

[25] Our focus on the present climate commitment leads to one particular value of R_{∞} . Of course, in any practical scenario, emissions will continue and R_{∞} will grow. In turn, the transient and ultimate climate commitments will increase and become more uncertain. Inasmuch as a substantially improved understanding of the role of aerosols in climate remains elusive, so will our ability to constrain future climate. In order to rule out the possibility that we already face a disturbingly large climate commitment, we need to rule out the possibility that the observed climate change has been driven by a climate forcing at the lower end of the range that is currently permitted by observations.

[26] **Acknowledgments.** We thank M. B. Baker, D. Battisti, C. M. Bitz, E. Steig, and L. Thompson for insightful discussions; two anonymous reviewers for constructive suggestions that greatly improved the manuscript; and Noah Diffenbaugh, the editor.

References

- Allen, M. R., et al. (2007), Observational constraints on climate sensitivity, in *Avoiding Dangerous Climate Change*, edited by H. J. Schellnhuber et al., pp. 281–289, Cambridge Univ. Press, Cambridge, U. K.
- Archer, D. (2005), Fate of fossil fuel CO_2 in geologic time, *J. Geophys. Res.*, *110*, C09S05, doi:10.1029/2004JC002625.
- Baker, M. B., and G. H. Roe (2009), The shape of things to come: Why is climate change so predictable?, *J. Clim.*, *22*, 4574–4589.
- Edwards, T. L., et al. (2007), Using the past to constrain the future: How the palaeorecord can improve estimates of global warming, *Prog. Phys. Geogr.*, *31*, 481–500.
- Forster, P., et al. (2007), Changes in atmospheric constituents and in radiative forcing, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. Solomon et al., pp. 129–234, Cambridge Univ. Press, Cambridge, U. K.
- Frame, D. J., B. B. Booth, J. A. Kettleborough, D. A. Stainforth, J. M. Gregory, M. Collins, and M. R. Allen (2005), Constraining climate forecasts: The role of prior assumptions, *Geophys. Res. Lett.*, *32*, L09702, doi:10.1029/2004GL022241.

- Gregory, J. M. (2000), Vertical heat transports in the ocean and their effect on time-dependent climate change, *Clim. Dyn.*, *16*, 501–515.
- Hansen, J., et al. (2005), Earth's energy imbalance: Confirmation and implications, *Science*, *308*, 1431–1435.
- Hare, B., and M. Meinshausen (2006), How much warming are we committed to and how much can be avoided?, *Clim. Change*, *75*, 111–149.
- Hegerl, G. C., et al. (2007), Understanding and attributing climate change, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. Solomon et al., pp. 663–745, Cambridge Univ. Press, Cambridge, U. K.
- Hoffert, M. I., A. J. Callegari, and C.-T. Hsieh (1980), The role of deep sea heat storage in the secular response to climatic forcing, *J. Geophys. Res.*, *85*, 6667–6679.
- Kiehl, J. T. (2007), Twentieth century climate model response and climate sensitivity, *Geophys. Res. Lett.*, *34*, L22710, doi:10.1029/2007GL031383.
- Knutti, R. (2008), Why are climate models reproducing the observed global surface warming so well?, *Geophys. Res. Lett.*, *35*, L18704, doi:10.1029/2008GL034932.
- Knutti, R. (2010), The end of model democracy?, *Clim. Change*, *102*, 395–404, doi:10.1007/s10584-010-9800-2.
- Knutti, R., and G. C. Hegerl (2008), The equilibrium sensitivity of the Earth's temperature to radiation changes, *Nat. Geosci.*, *1*, 735–743.
- Lyman, J. M., et al. (2010), Robust warming of the global upper ocean, *Nature*, *465*, 334–337.
- Matthews, H. D., and A. J. Weaver (2010), Committed climate warming, *Nat. Geosci.*, *3*, 142–143.
- Meehl, G. A., et al. (2005), How much more global warming and sea level rise?, *Science*, *307*, 1769–1772.
- Myhre, G., E. J. Highwood, K. P. Shine, and F. Stordal (1998), New estimates of radiative forcing due to well mixed greenhouse gases, *Geophys. Res. Lett.*, *25*, 2715–2718.
- Plattner, G.-K., et al. (2008), Long-term climate commitments projected with climate-carbon cycle models, *J. Clim.*, *21*, 2721–2751.
- Purkey, S. G., and G. C. Johnson (2010), Warming of global abyssal and deep southern ocean waters between the 1990s and 2000s: Contributions to global heat and sea level rise budgets, *J. Clim.*, doi:10.1175/2010JCLI3682.1, in press.
- Ramanathan, V., and Y. Feng (2008), On avoiding dangerous anthropogenic interference with the climate system: Formidable challenges ahead, *Proc. Natl. Acad. Sci. U. S. A.*, *38*, 14,245–14,250.
- Raper, S. C. B., et al. (2001), Use of an upwelling-diffusion energy balance climate model to simulate and diagnose A/OGCM results, *Clim. Dyn.*, *17*, 601–613.
- Schwartz, S. E., et al. (2007), Quantifying climate change—Too rosy a picture?, *Nat. Rep. Clim. Change*, *2*, 23–24.
- Solomon, S., et al. (2009), Irreversible climate change due to carbon dioxide emissions, *Proc. Natl. Acad. Sci. U. S. A.*, *106*, 1704–1709.
- Trenberth, K. E., et al. (2007), Observations: Surface and atmospheric climate change, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. Solomon et al., pp. 235–336, Cambridge Univ. Press, Cambridge, U. K.
- Wigley, T. M. L. (2005), The climate change commitment, *Science*, *307*, 1766–1769.
- Winton, M., et al. (2009), Importance of ocean heat uptake efficacy to transient climate change, *J. Clim.*, *23*, 2333–2344.

K. C. Armour, Department of Physics, University of Washington, Box 351560, Seattle, WA 98195, USA. (karmour@u.washington.edu)
 G. H. Roe, Department of Earth and Space Sciences, University of Washington, Seattle, WA 98195, USA. (gerard@ess.washington.edu)